

Test of the conserved vector current hypothesis in the beta-decay of ^{14}C

I. Ahmad[‡], S. J. Freedman^{*†}, B. K. Fujikawa^{*}, J. P. Greene[‡], J. L. Mortara^{*†}, J. P. Schiffer[‡], and A. R. Zeuli[‡]

In 1957, Feynman and Gell-Mann proposed the conserved vector current (CVC) hypothesis[1]. In the classic case of the mass-12 isospin triplet (^{12}B - ^{12}C - ^{12}N), CVC relates the width of the analog electromagnetic transition ($^{12}\text{C}^* \rightarrow ^{12}\text{C}$) to the shape factors of the beta decay spectra ($^{12}\text{B} \rightarrow ^{12}\text{C}$) and ($^{12}\text{N} \rightarrow ^{12}\text{C}$). The shape factor represents a deviation of the beta spectrum from the simple allowed shape by the additional factor $S(E)=1+a E$, where E is the total electron energy. Several experiments have measured the shape factor in the mass-12 system, but the agreement with CVC is weak at best[2]. The mass-14 isospin triplet (^{14}C - ^{14}N - ^{14}O) represents another viable system for testing CVC. We have performed a measurement of the beta decay spectrum of $^{14}\text{C} \rightarrow ^{14}\text{N}$ (Q 156 keV) toward this end. The apparatus used for this measurement consists of a superconducting solenoid and a Si(Li) solid state detector[3]. The magnetic field transports the electrons in helical orbits ($r < 3$ mm) to the Si(Li) detector without the possibility of scattering on material collimators. The response of the detector was determined by measurement of internal conversion spectra from ^{139}Ce and ^{109}Cd . The ^{14}C data consists of four separate runs with a total of about 7×10^9 total decays accumulated over a period of 515 live hours. The data is fitted from 65 to 250 keV. The results are quoted for this energy interval, but the shape factor was observed to be independent of the chosen interval. In Fig. 1 we show the fit from the first run which yielded shape factor of $a = -(39.1 \pm 0.4) \times 10^{-2}$ per MeV. Combining this result with three additional data runs, we have shape factor of $a = -(39.1 \pm 0.2_{\text{stat}} \pm 0.6_{\text{syst}}) \times 10^{-2}$ per MeV. This represents the most precise determination of the shape factor in ^{14}C and appears to be in good agreement with the value predicted by CVC of $a = -(38.0 \pm 1.2) \times 10^{-2}$ per MeV where the error reflects the uncertainty in the

radiative width of $^{14}\text{C} \rightarrow ^{14}\text{N}^*$. This result will be complemented by a measurement of the ^{14}O shape factor[4].

Footnotes and References

*Lawrence Berkeley National Laboratory

†University of California, Berkeley

‡Argonne National Laboratory

1. R. P. Feynman and M. Gell-Mann, Phys. Rev. **109**, 193 (1957).
2. F. P. Calaprice and B. R. Holstein, Nucl. Phys. **A273**, 301 (1976).
3. J. L. Mortara *et al.*, Phys. Rev. Lett. **70**, 394 (1993).
4. J. T. Burke, *et al.*, 1999 NSD Annual Report.




Fig. 1. Top: the Mass-14 isospin triplet. Bottom: residuals of fit to ^{14}C spectrum.